

Appl. No. 09/744,267  
 Docket No. CM1882  
 Amdt. Dated March 23, 2009  
 Reply to Office Action Dated December 23, 2008  
 Customer No. 27752

RECEIVED  
 CENTRAL FAX CENTER


MAR 23 2009

### AMENDMENTS TO THE CLAIMS

This listing of claims will replace all prior versions, and listings, of claims in the application:

1. (Currently amended) A fabric care composition comprising:

- i)- from 5% to 10% of a nitrogen containing dye fixing agent,
- ii)- from 0.001 % to 20 % of a scum reducing agent comprising a polyoxyalkylene alkyl amine surface active agent having the formula




Wherein R is selected from C7-C21 linear alkyl, C7-C21 branched alkyl, C7-C21 linear alkenyl, C7-C21 branched alkenyl, and mixtures thereof, R<sup>1</sup> is ethylene, R<sup>2</sup> is selected from C3-C4 linear alkyl, C3-C4 branched alkyl, 1, 2 propylene, and mixtures thereof, and

- iii)- a polyamino-functional polymer wherein said polymer comprises a polyamine backbone corresponding to the formula:



having a polyamine formula  $\text{V}_{(n+1)}\text{W}_m\text{Y}_n\text{Z}$  or a polyamine backbone corresponding to the formula:



having a polyamine formula  $\text{V}_{(n-k+1)}\text{W}_m\text{Y}_n\text{Y}'_k\text{Z}$ , wherein k is less than or equal to n, said polyamine backbone has a molecular weight greater than 200 daltons, wherein

- i) V units are terminal units having the formula:


Appl. No. 09/744,267

Docket No. CM1882


Amdt. Dated March 23, 2009

Reply to Office Action Dated December 23, 2008


Customer No. 27752




ii) W units are backbone units having the formula:




iii) Y units are branching units having the formula:



iv) Y' units are branch point for a backbone or branch ring having the formula:



v) Z units are terminal units having the formula:



wherein backbone linking R units are selected from the group consisting of C<sub>2</sub>-C<sub>12</sub> alkylene, C<sub>4</sub>-C<sub>12</sub> alkenylene, C<sub>3</sub>-C<sub>12</sub> hydroxyalkylene, C<sub>4</sub>-C<sub>12</sub> dihydroxy-alkylene, C<sub>8</sub>-C<sub>12</sub> dialkylarylene, -(R<sup>1</sup>O)<sub>x</sub>R<sup>1</sup>-, -(R<sup>1</sup>O)<sub>x</sub>R<sup>5</sup>(OR<sup>1</sup>)<sub>x</sub>-, -(CH<sub>2</sub>CH(OR<sup>2</sup>)CH<sub>2</sub>O)<sub>z</sub>(R<sup>1</sup>O)<sub>y</sub>R<sup>1</sup>(OCH<sub>2</sub>CH(OR<sup>2</sup>)CH<sub>2</sub>)<sub>w</sub>-, -C(O)(R<sup>4</sup>)<sub>t</sub>C(O)-, -CH<sub>2</sub>CH(OR<sup>2</sup>)CH<sub>2</sub>-, and mixtures thereof; wherein R<sup>1</sup> is selected from the group consisting of C<sub>2</sub>-C<sub>6</sub> alkylene and mixtures thereof; R<sup>2</sup> is selected from the group consisting of hydrogen, -(R<sup>1</sup>O)<sub>x</sub>B, and mixtures thereof; R<sup>4</sup> is selected from the group consisting of C<sub>1</sub>-C<sub>12</sub> alkylene, C<sub>4</sub>-C<sub>12</sub> alkenylene, C<sub>8</sub>-C<sub>12</sub> arylalkylene, C<sub>6</sub>-C<sub>10</sub> arylene, and mixtures thereof; R<sup>5</sup> is selected from the group consisting of C<sub>1</sub>-C<sub>12</sub> alkylene, C<sub>3</sub>-C<sub>12</sub> hydroxyalkylene, C<sub>4</sub>-C<sub>12</sub> dihydroxy-alkylene, C<sub>8</sub>-C<sub>12</sub> dialkylarylene, -C(O)-, -C(O)NHR<sup>6</sup>NHC(O)-, -R<sup>1</sup>(OR<sup>1</sup>)-, -C(O)(R<sup>4</sup>)<sub>t</sub>C(O)-, -CH<sub>2</sub>CH(OH)CH<sub>2</sub>-, -CH<sub>2</sub>CH(OH)CH<sub>2</sub>O(R<sup>1</sup>O)<sub>y</sub>R<sup>1</sup>OCH<sub>2</sub>CH(OH)CH<sub>2</sub>-, and mixtures thereof; R<sup>6</sup> is selected


Appl. No. 09/744,267  
 Docket No. CM1882  
 Amdt. Dated March 23, 2009  
 Reply to Office Action Dated December 23, 2008  
 Customer No. 27752

from the group consisting of C<sub>2</sub>-C<sub>12</sub> alkylene or C<sub>6</sub>-C<sub>12</sub> arylene; R' units are selected from the group consisting of hydrogen, C<sub>1</sub>-C<sub>22</sub> alkyl, C<sub>3</sub>-C<sub>22</sub> alkenyl, C<sub>7</sub>-C<sub>22</sub> arylalkyl, C<sub>2</sub>-C<sub>22</sub> hydroxyalkyl, -(CH<sub>2</sub>)<sub>p</sub>CO<sub>2</sub>M, -(CH<sub>2</sub>)<sub>q</sub>SO<sub>3</sub>M, -CH(CH<sub>2</sub>CO<sub>2</sub>M)CO<sub>2</sub>M, -(CH<sub>2</sub>)<sub>p</sub>PO<sub>3</sub>M, -(R<sup>1</sup>O)<sub>x</sub>B, -C(O)R<sup>3</sup>, and mixtures thereof; B is selected from the group consisting of hydrogen, C<sub>1</sub>-C<sub>6</sub> alkyl, -(CH<sub>2</sub>)<sub>q</sub>SO<sub>3</sub>M, -(CH<sub>2</sub>)<sub>p</sub>CO<sub>2</sub>M, -(CH<sub>2</sub>)<sub>q</sub>(CHSO<sub>3</sub>M)CH<sub>2</sub>SO<sub>3</sub>M, -(CH<sub>2</sub>)<sub>q</sub>-(CHSO<sub>2</sub>M)CH<sub>2</sub>SO<sub>3</sub>M, -(CH<sub>2</sub>)<sub>p</sub>PO<sub>3</sub>M, -PO<sub>3</sub>M, and mixtures thereof; R<sup>3</sup> is selected from the group consisting of C<sub>1</sub>-C<sub>18</sub> alkyl, C<sub>7</sub>-C<sub>12</sub> arylalkyl, C<sub>7</sub>-C<sub>12</sub> alkyl substituted aryl, C<sub>6</sub>-C<sub>12</sub> aryl, and mixtures thereof; M is hydrogen or a water soluble cation in sufficient amount to satisfy charge balance; X is a water soluble anion; m has the value from 2 to 700; n has the value from 0 to 350; p has the value from 1 to 6, q has the value from 0 to 6; r has the value of 0 or 1; w has the value 0 or 1; x has the value from 1 to 100; y has the value from 0 to 100; z has the value 0 or 1, wherein the weight ratio of scum reducing agent to the sum of the polyamino-functional polymer and the dye fixing agents is from 0.05:1 to 2:1.


2. (Cancelled)

3. (Cancelled)

4. (Previously Presented) A composition according to Claim 1, wherein the polyoxyalkylene alkyl amine surface active agent has the formula:



wherein R is selected from C<sub>7</sub>-C<sub>21</sub> linear alkyl, C<sub>7</sub>-C<sub>21</sub> branched alkyl, C<sub>7</sub>-C<sub>21</sub> linear alkenyl, C<sub>7</sub>-C<sub>21</sub> branched alkenyl, and mixtures thereof; R<sup>1</sup> is ethylene; R<sup>2</sup> is selected from C<sub>3</sub>-C<sub>4</sub> linear alkyl, C<sub>3</sub>-C<sub>4</sub> branched alkyl, and mixtures thereof; R<sup>3</sup> is selected from hydrogen, C<sub>1</sub>-C<sub>4</sub> linear alkyl, C<sub>3</sub>-C<sub>4</sub> branched alkyl, and mixtures thereof; R<sup>4</sup> is selected from hydrogen, C<sub>1</sub>-C<sub>4</sub> linear alkyl, C<sub>3</sub>-C<sub>4</sub> branched alkyl, and mixtures thereof; A is



Appl. No. 09/744,267  
Docket No. CM1882  
Amdt. Dated March 23, 2009  
Reply to Office Action Dated December 23, 2008  
Customer No. 27752

$R^5$  is selected from  $-(R^1O)_x(R^2O)_y$  unit,  $C_1$ - $C_{16}$  linear alkyl,  $C_1$ - $C_{16}$  branched alkyl,  $C_1$ - $C_{16}$  linear alkenyl,  $C_1$ - $C_{16}$  branched alkenyl, and mixtures thereof;  
wherein the index  $m$  is 1 or 2, the index  $n$  is 0 or 1, provided that when  $m$  is equal to 1,  $n$  is equal to 1; and when  $m$  is 2  $n$  is 0;  
wherein the index  $x$  is from 0 to about 50, preferably from 1 to 25,  
wherein the index  $y$  is from 0 to about 10;  
wherein the index  $q$  is 0 or 1.

5. (Previously Presented) A composition according to Claim 4, wherein said index  $x$  is from 1 to 25.

6. (Previously Presented) A composition according to Claim 5, wherein said index  $m$  is equal to 2 and  $n$  is equal to 0.

7. (Cancelled)

8. (Previously Presented) A composition according to Claim 7, wherein said dye fixing agent is a cellulose reactive dye fixing agent.

9.-14. (Cancelled)

15. (Previously Presented) A composition according to Claim 1, wherein the weight ratio of the scum reducing agent to the sum of the polyamino-functional polymer and dye fixing agent is from 0.1:1 to 1:1.

16. (Previously Presented) A composition according to Claim 1, further comprising an ease of formulation solvent having a ClogP of from about 0.15 to about 0.64.

17. (Previously Presented) A composition according to Claim 16, wherein the ease of formulation solvent is selected from the group consisting of: mono-ols, C6 diols, C7 diols, octanediol isomers, butanediol derivatives, trimethylpentanediol isomers, ethylmethylpentanediol isomers, propyl pentanediol isomers, dimethylhexanediol isomers, ethylhexanediol isomers, methylheptanediol isomers, octanediol isomers, nonanediol

Appl. No. 09/744,267  
Docket No. CM1882  
Amdt. Dated March 23, 2009  
Reply to Office Action Dated December 23, 2008  
Customer No. 27752

isomers, alkyl glyceryl ethers, di(hydroxy alkyl) ethers, and aryl glyceryl ethers, aromatic glyceryl ethers, alicyclic diols and derivatives, C3C7 diol alkoxyated derivatives, aromatic diols, and unsaturated diols.

18. (Previously Presented) A composition according to Claim 16, wherein the ease of formulation solvent is selected from the group consisting of 1,2-Hexanediol, 2-Ethyl-1,3-hexanediol and 2,2,4-Trimethyl-1,3-pentanediol.

19. (Previously Presented) A composition according to Claim 16, wherein said ease of formulation solvent comprises an asymmetric solvent.

20. (Previously Presented) A composition according to Claim 19, wherein the composition is essentially clear.

21. (New) A composition according to Claim 1, wherein the R<sup>2</sup> of the polyoxyalkylene alkyl amine surface active agent is a 1,2-prolylene.

22. (New) A composition according to Claim 1, wherein the polyoxyalkylene alkyl amine surface active agent comprises a ratio of R<sup>1</sup> to R<sup>2</sup> from about 4 to about 12 ethylene units to about 1 to about 4 1,2-prolylene units.